



Brachyglottis repanda

Common Name(s):

rangiora, bushman's toilet paper, bushman's friend

Current Threat Status (2012):

Not Threatened

Distribution:

Endemic. North Island throughout. South Island - north west Nelson to just south of Greymouth in the west, and near Kekerengu in the east. Naturalised on Banks Peninsula, Otago Peninsula, and on Stewart Island at Oban.

Habitat:

Common in coastal, lowland and lower montane shrubland and open forest. Often a pioneer species.

Features:

Shrub to small tree up to 6 m or more tall. Trunk one or more arising from ground, covered in somewhat corky bark. Branches stout, spreading, rather brittle, initially densely clad in fine white to buff tomentum becoming glabrescent with age. Petiole stout, grooved, 80-100 mm long. Leaves leathery, 50-250(-300) X 50-20(-30) mm, dark green to pale green above, undersides clad in fine, appressed vivid white hairs, broad- to ovate-oblong, obtuse to subacute, obliquely cordate to truncate at base, margins distantly dentately lobed to sinuate. Inflorescence a much branched panicle. Capitula 5 mm diam., numerous, without ligules (discoïd). Involucral bracts 3 mm long, narrow-oblong to narrow spatulate, margins scarious except at base. Florets 10-12, yellow. Seeds (cypsela) narrowly oblong-elliptic to oblong elliptic, 1-1.8 mm long, ribs 6, rounded, broad. Pappus 2-3 mm, buff-yellow, scabrid.

Flowering:

(July-) August-October
(-November)

Fruiting:

(October-) November-
December (-January)

Threats:

Not Threatened

References and further reading:

Thorsen, M. J.; Dickinson, K. J. M.; Seddon, P. J. 2009. Seed dispersal systems in the New Zealand flora. Perspectives in Plant Ecology, Evolution and Systematics 2009 Vol. 11 No. 4 pp. 285-309

For more information, visit:

http://nzpcn.org.nz/flora_details.asp?ID=1562



Caption: *Brachyglottis repanda*
Photographer: Wayne Bennett



Caption: *Brachyglottis repanda*
Photographer: Wayne Bennett