

Hymenophyllum dilatatum

COMMON NAME

filmy fern, matua mauku

SYNONYMS

Mecodium dilatatum (G. Forst.) Copel.; Sphaerocionium dilatatum (G.Forst.) C.Presl; Trichomanes dilatatum G.Forst.; Diploophyllum dilatatum (G.Forst.) Bosch; Leptocionium sororium C.Presl

FAMILY

Hymenophyllaceae

AUTHORITY

Hymenophyllum dilatatum (G. Forst.) Sw.

FLORA CATEGORY

Vascular – Native

ENDEMIC TAXON

Yes

ENDEMIC GENUS

No

ENDEMIC FAMILY

No

STRUCTURAL CLASS

Ferns

NVS CODE

HYMDIL

CHROMOSOME NUMBER

2n = 72

CURRENT CONSERVATION STATUS

2017 | Not Threatened

PREVIOUS CONSERVATION STATUSES

2012 | Not Threatened

2009 | Not Threatened

2004 | Not Threatened

DISTRIBUTION

Endemic, New Zealand: North Island, South Island, Stewart Island/Rakiura, Chatham Islands, Auckland Islands. Widespread except for the drier parts of the eastern South Island.

HABITAT

Coastal to montane in forest. Usually epiphytic or on fallen logs and banks, Very rarely on the forest floor or on boulders.


WETLAND PLANT INDICATOR STATUS RATING

UPL: Obligate Upland

Rarely is a hydrophyte, almost always in uplands (non-wetlands).



Coromandel, June. Photographer: John Smith-Dodsworth, Licence: CC BY-NC.



Coromandel, June. Photographer: John Smith-Dodsworth, Licence: CC BY-NC.

DETAILED DESCRIPTION

Epiphytic (very rarely terrestrial) fern. **Rhizomes** long-creeping, gracile, wiry when fresh very brittle when dry. **Stipes** often widely spaced on rhizomes, 20–150–(200) mm long, stout, glabrous, distinctly though narrowly winged for part of length; rachises broadly winged throughout. **Laminae** 80–400–(800) × 40–150–(160) mm, ovate, narrowly ovate to lanceolate, 3–4-pinnate, bright to dark green, glabrous. **Ultimate segments** rather broad, margins smooth, plane. **Sori** terminating ultimately segments, slightly sunk in lamina, many on each primary pinna. **Indusial flaps** smooth. (Description adapted from Brownsey & Smith-Dodsworth (2000)).

SIMILAR TAXA

Easily distinguished from all other *Hymenophyllum* by the glabrous, smooth margined fronds; by the conspicuous broad, flat wing bordering much of the stipe and all of the rachis; and by the very broad pinnae.

LIFE CYCLE

Minute spores are wind dispersed (Thorsen et al., 2009).

PROPAGATION TECHNIQUE

Difficult—should not be removed from the wild

ETYMOLOGY

hymenophyllum: Membranous leaf, from the Greek humen and phullon

WHERE TO BUY

Not commercially available

ATTRIBUTION

Fact sheet prepared for NZPCN by P.J. de Lange (17 April 2011). Description adapted from Brownsey & Smith-Dodsworth (2000).

REFERENCES AND FURTHER READING

Brownsey PJ, Smith-Dodsworth JC. 2000. New Zealand Ferns and Allied Plants. David Bateman, Auckland, NZ. 168 p.

Thorsen MJ, Dickinson KJM, Seddon PJ. 2009. Seed dispersal systems in the New Zealand flora. *Perspectives in Plant Ecology, Evolution and Systematics* 11: 285–309. <https://doi.org/10.1016/j.ppees.2009.06.001>.

NZPCN FACT SHEET CITATION

Please cite as: de Lange, P.J. (Year at time of access): *Hymenophyllum dilatatum* Fact Sheet (content continuously updated). New Zealand Plant Conservation Network.

<https://www.nzpcn.org.nz/flora/species/hymenophyllum-dilatatum/> (Date website was queried)

MORE INFORMATION

<https://www.nzpcn.org.nz/flora/species/hymenophyllum-dilatatum/>