

12. Plant identification


There are many sources of information to help identify plants:

1. **Written and pictorial descriptions** of plants, which you can compare with your unknown specimen to aid in its identification. Good descriptions direct you to crucial diagnostic features for the relevant taxon, explain the range variability found and point out botanical and ecological characteristics of importance. A good source of information about native plants is the website of the New Zealand Plant Conservation Network – see www.nzpcn.org.nz. This site has information (including photos) about all New Zealand’s indigenous and naturalised exotic plants.

Please note—to use an identification guide properly, you need to know enough of the vocabulary to understand the choices presented to you. All good identification guides provide a glossary and a list of abbreviations to help with this.

2. **Ask an expert.** Many people around the country, including professional botanists and members of botanical societies, may be able to assist with identification. Providing fresh samples will help accurate identification, but may not always be possible. Ensure that you have the appropriate permission before collecting specimens (see page 59). Alternatively, photographs that show significant features can be used. A desktop scanner is a useful way to make an image of the specimen if you don’t have a good camera.

3. **Keys** help you find the likely description of your specimen rapidly and simply. Most keys are arranged to present you with a series of choices (decision points), usually dichotomous (dividing in two). The paired statements of each ‘couplet’ are framed to be contrasting and mutually exclusive. Each choice you make narrows down the possibilities for your specimen until you find the appropriate description. Terminology is precise and brief. Keys can vary in complexity. An easy one to use is *Which Native Tree?* by Andrew Crowe.



Advice for using keys

- Note down the route taken—to trace path in case you need to back track
- Read full description of both choices for each step
- Do not guess—consult glossary if precise meaning of term unknown. Where measurements are required, use a ruler
- If features are very small, use an appropriate lens to inspect them clearly.
- If key is multi-part one, look carefully at the descriptions for higher levels of taxa before progressing to species key.
- If both pairs of choices seem reasonable, try each route—one will usually prove to be unsuitable at a later stage.

You may like to try out the following key on a kōwhai *Sophora* sp. specimen:



A selection of *Sophora*: Clockwise from above: *S. chathamica* (photo: Geoff Walls), *S. fulvida*, *S. molloyi*, *S. prostrata*.


Sophora (FABACEAE) in New Zealand: taxonomy, distribution, and biogeography
P. B. Heenan, P.J. de Lange and A.D. Wilton

Key to the New Zealand kōwhai species (adapted from *NZ Journal Of Botany* 2001, Volume 39)


This key is only for New Zealand species of *Sophora*, and does not include hybrid material. If the key does not work, the plant material should be checked to see if it is hybrid origin.

- 1 Shrub, branches at or near ground level and usually slender2
Tree, branches well above ground level and usually thick 5
- 2 Branchlets not interlaced, usually grey to grey-brown, glabrous to sparsely hairy..... 3
Branchlets interlaced, usually yellow-brown to orange-brown, sparsely to moderately hairy..... 4
- 3 Shrub usually wider than high; main branches spreading to decumbent, sometimes prostrate,
underground branches and rhizomes absent; leaves 23–37 leaflets; Leaflets 5.0–12.0 × 2.0–6.0 mm
elliptic, elliptic oblong, to broadly elliptic, sparsely hairy; Kapiti island, islands in Cook Strait,
southern headlands of North Island, dry and exposed windy bluffs ***S. molloyi***
Shrub usually of similar width and height; main branches upright to spreading, underground
branches and rhizomes usually present, often with numerous branches near the base; leaves
with 35–52 leaflets; leaflets 3.3–5.8 × 2.5–3.1 mm, orbicular, obovate, to oblong obovate , usually
more or less glabrous; northern Nelson, western Marlborough, marble and limestone outcrops
.....***S. longicarinata***
- 4 Branchlets interlaced on juvenile and adult; leaves <3 cm long, leaflet pairs 1–5, usually
glabrous; standard petal orange; pods lacking wings; seeds dark brown; eastern South Island,
dry grey scrub communities ***S. prostrata***
Branchlets interlaced in juvenile only; leaves > 3cm, leaflet pairs > sparsely to moderately hairy;
Flowers usually absent, if present standard petal yellow; pods usually absent, if present
winged; seeds yellow or yellow brown; North and South Islands, terraces and hillslopes..... ***S. microphylla***
5. Ovary and leaves with hairs spreading, curved and/or twisted; leaflets densely hairy 6
Ovary and leaves with hairs appressed, straight, leaflets not densely hairy, or if densely
hairy greater than 15 mm long 7
- 6 Leaves with 61–91 leaflets; leaflets elliptic to elliptic oblong, occasionally narrowly obovate
usually sessile; leaflet hairs appressed, decumbent, or spreading, predominantly straight
and sometimes twisted; northern North Island, andesitic and volcanic rock outcrops..... ***S. fulvida***
Leaves with 47–75 leaflets; leaflets ovate, broadly elliptic, to sometimes more or less
orbicular, with a more or less petiolule; leaflet hairs appressed, decumbent, spreading or
patent, predominantly curly, curved, or twisted, central North Island, siltstone, sandstone
and mudstone (papa).....***S. godleyi***
- 7 Leaves with fewer than 23 leaflets; leaflets more than 18 mm long, three times longer than
wide narrowly elliptic to elliptic-oblong, densely hairy; eastern North Island, terraces and
hillslopes***S. tetraptera***
Leaves with more than 24 leaflets; leaflets less than 16 mm long, length usually less than
twice their width, elliptic, broadly elliptic, obovate, broadly obovate, ovate, broadly ovate,
oblong to more or less orbicular, glabrous or moderately hairy 8
- 8 Juvenile growth present, leaflets 4.5–12.5 × 2.3–5.7 mm, distal and proximal leaflets
usually similar in size, distant, not crowded or overlapping , elliptic, broadly elliptic,
obovate to ovate, sometimes more or less orbicular, usually moderately hairy; North
and South Islands, terraces and hill country ***S. microphylla***
Juvenile growth habit absent, leaflets 6.0–16.0 × 4.0–8.0 mm, distal leaflets usually
smaller than proximal leaflets, crowded and overlapping, broadly elliptic, broadly
obovate, obovate to more or less orbicular, moderately hairy; North Island and Chatham
Islands, coastal and lowland hill country.....***S. chathamica***
Juvenile growth absent; leaflets 3.3–5.8 × 2.5–3.1 mm, distal and proximal leaflets
similar in size, overlapping to distant, orbicular, obovate, to oblong-obovate, more or
less glabrous; Northern Nelson, western Marlborough, marble and Limestone rock
outcrops***S. longicarinata***

9. (a) Bark very thick and stringy. Large forest tree. Seed ovoid. ***Podocarpus totara***
tōtara (40 m)
PODOCARPACEAE
- (b) Bark thinner more papery and flaky. Leaves often larger than true tōtara esp. ***Podocarpus cunninghamii***
(formerly known as *P. hallii*), thin-
barked tōtara, Hall's tōtara (20 m)
PODOCARPACEAE
- (c) Bark thin and stringy, mature leaves greater than 10 mm long, needle-leaved ***Podocarpus acutifolius***
prickly tōtara (9 m)
PODOCARPACEAE
- (d) Prostrate to low growing alpine shrub, branches wide-spreading, leaves up ***Podocarpus nivalis***
to 15mm long. snow tōtara (5 m)
PODOCARPACEAE
10. (a) Seeds with black flesh (like little plum). Leaves bluish green above and
whitish below. in two irregular rows on branchlets. Leaf tips are rounded with
a small sharp point. Bark bluish, almost black, shiny, hammer marked; loose
flakes leave red marks on trunk when shed. Juvenile has untidy interlacing ***Prumnopitys taxifolia***
slender branchlets with a mixture of scale like leaves and larger linear leaves,
bronze-green. matai, black pine (40 m)
PODOCARPACEAE
- (b) Seeds with red flesh (like little plum). Spirally arranged leaves are distinctly
flattened into 2 rows, narrowed to a point and green underneath. Juvenile ***Pectinopitys ferruginea***
and adolescent leaves 1–3 cm long. Bark dark grey to grey brown, rough also
hammer marked but not leaving red marks as in matai. miro, brown pine (25 m)
PODOCARPACEAE
11. (a) Trees with black spherical seed on fleshy orange stalks. Juvenile with leaves
in two rows, and branches not weeping, adult leaves are 2–3 mm long, awl- ***Dacrycarpus dacrydioides***
like closely appressed to the branchlets. Branchlets are characteristically
upturned at their tips. Bark grey and "hammer marked". Buttress roots. kahikatea/ white pine (65 m)
PODOCARPACEAE
- (b) Trees with black pear-shaped seeds projecting from red fleshy cups. Juvenile ***Dacrydium cupressinum***
leaves not in distinct row, branches weeping, bark dark brown and coming
off in large flakes. rimu, red pine (60 m)
PODOCARPACEAE
12. (a) Trees with ribbed seeds seated on orange fleshy cushions. Juvenile foliage
large 15–40 mm long linear leaves. Abrupt transition to adult foliage ***Halocarpus kirkii***
consisting of 2–3 mm long, overlapping scale leaves, weakly keeled on back. Northern North island only. manaoa, Kirk's rimu (25 m)
PODOCARPACEAE
- (b) Shrub showing abrupt transition from leafy juvenile shoots bearing linear
leaves to narrow adult shoots with appressed scale leaves Juvenile leaves ***Halocarpus bidwillii***
5–10 mm long, 1–1.5 mm wide; fleshy outgrowth at base of ribbed seed
white-yellow. bog pine (5 m)
PODOCARPACEAE
- (c) Erect small tree, bark scaling off in small flakes, reddish silver beneath. ***Halocarpus biformis***
Juvenile leaves 1–20 mm long; adult leaves with prominent keel on back. pink pine (9 m)
Fleshy outgrowth at base of ribbed seed orange (as for *H. kirkii*).
PODOCARPACEAE
- (d) Rounded bush or small tree with seeds half hidden in surrounding greenish
fleshy scales. Bark thin grey to reddish brown not flaking, unfissured. Juvenile ***Manoao colensoi***
foliage on erect or spreading shoots. Adult leaves strongly keeled, shoot tips
erect, seed subtended and half hidden by green to whitish green fleshy scale. manaoa, silver pine (15 m)
PODOCARPACEAE
- (e) Small tree, bark thin, grey, Juvenile foliage on drooping shoots; leaves of
semi-adult stage bristling all round the shoot; adult scale leaves weakly ***Lepidothamnus intermedius***
keeled; shoot tips curving over and down. 2–3 yellow fleshy scale leaves
subtending seed. yellow-silver pine (15 m)
PODOCARPACEAE




Kauri
(*Agathis australis*)



Kawaka
(*Libocedrus plumosa*)



Kaikawaka
(*Libocedrus bidwillii*)



Tānekaha
(*Phyllocladus trichomanoides*)




Toatoa
(*Phyllocladus toatoa*)




Alpine toatoa
(*Phyllocladus alpinus*)



Prickly tōtara
(*Podocarpus acutifolius*)




Thin-barked tōtara
(*Podocarpus cunninghamii*)




Tōtara
(*Podocarpus totara*)



Snow tōtara
(*Podocarpus nivalis*)



Matai
(*Prumnopitys taxifolia*)



Miro
(*Prumnopitys ferruginea*)



Kahikatea
(*Dacrycarpus dacrydioides*)




Rimu
(*Dacrydium cupressinum*)




Manoao
(*Halocarpus kirkii*)



Bog pine
(*Halocarpus bidwillii*)




Pink pine
(*Halocarpus biformis*)



Manoao / silver pine
(*Manoao colensoi*)



Yellow silver pine
(*Lepidothamnus intermedius*)



Pigmy pine
(*Lepidothamnus laxifolius*)

N.M.ADAMS

Gymnosperm illustrations from *Trees and Shrubs of New Zealand* by AL Poole and Nancy M Adams, © Landcare Research New Zealand 1994.

